

# 6 Fuzzy Relations

*Fuzzy Systems Engineering  
Toward Human-Centric Computing*

# Contents

**6.1 The concept of relations**

**6.2 Fuzzy relations**

**6.3 Properties of fuzzy relations**

**6.4 Operations on fuzzy relations**

**6.5 Cartesian product, projections, and cylindrical extension**

**6.6 Reconstruction of fuzzy relations**

**6.7 Binary fuzzy relations**


# 6.1 The concept of relations

# Relation

Docs

**X**

$\{d_1, d_2, \dots, d_i, \dots, d_n\}$




$d_i$

Keywords

**Y**



$\{w_1, w_2, \dots, w_j, \dots, w_m\}$



$w_j$

$$R = \{(d_i, w_j) \mid d_i \in \mathbf{X}, w_j \in \mathbf{Y}\}$$

Relation  $R : \mathbf{X} \times \mathbf{Y} \rightarrow \{0,1\}$




$\mathbf{X} = \mathbf{Y} = \{2, 4, 6, 8\}$   
equal to  
 $R = \{(2,2), (4,4), (6,6), (8,8)\}$

$$R = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$


# Examples

## Circle



$$R(x, y) = \begin{cases} 1 & \text{if } x^2 + y^2 = r^2 \\ 0 & \text{otherwise} \end{cases}$$

## Square



$$R(x, y) = \begin{cases} 1 & \text{if } |x| \leq 1 \text{ and } |y| \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

## 6.2 Fuzzy relations

# Fuzzy relation $R : \mathbf{X} \times \mathbf{Y} \rightarrow [0,1]$

## Example


Docs

$$\mathbf{D} = \{d_{fs}, d_{nf}, d_{ns}, d_{gf}\}$$

Keywords

$$\mathbf{W} = \{w_f, w_n, w_g\}$$

$$R : \mathbf{D} \times \mathbf{W} \rightarrow [0,1]$$




$$R = \begin{matrix} & \begin{matrix} w_f & w_n & w_g \end{matrix} \\ \begin{bmatrix} 1 & 0 & 0.6 \\ 0.8 & 1 & 0 \\ 0 & 1 & 0 \\ 0.8 & 0 & 1 \end{bmatrix} & \begin{matrix} d_{fs} \\ d_{nf} \\ d_{ns} \\ d_{gf} \end{matrix} \end{matrix}$$



# Example

$$R_e(x, y) = \exp\left\{\frac{-|x - y|}{\alpha}\right\}, \quad \alpha > 0$$

$$\mathbf{X} = \mathbf{Y} = [0, 4]$$



$x$  approximately equal to  $y$

$$\alpha = 1$$

## 6.3 Properties of fuzzy relations

Fuzzy relation  $R : \mathbf{X} \times \mathbf{Y} \rightarrow [0,1]$

Domain

$$\text{dom}R(x) = \sup_{y \in \mathbf{Y}} R(x, y)$$

Codomain

$$\text{cod}R(y) = \sup_{x \in \mathbf{X}} R(x, y)$$

# Representation of fuzzy relations

$$R = \bigcup_{\alpha \in [0,1]} \alpha R_{\alpha}$$

$$R(x, y) = \sup_{\alpha \in [0,1]} \{ \min[\alpha, R(x, y)] \}$$

Representation theorem

Fuzzy relations  $P, Q : \mathbf{X} \times \mathbf{Y} \rightarrow [0,1]$

Equality

$$P(x,y) = Q(x,y) \quad \forall (x,y) \in \mathbf{X} \times \mathbf{Y}$$

Inclusion

$$P(x,y) \leq Q(x,y) \quad \forall (x,y) \in \mathbf{X} \times \mathbf{Y}$$

# 6.4 Operations on fuzzy relations

Fuzzy relations  $P, Q : \mathbf{X} \times \mathbf{Y} \rightarrow [0,1]$

Union:  $R = P \cup Q$

$$R(x,y) = P(x,y) \textit{ s } Q(x,y) \quad \forall (x,y) \in \mathbf{X} \times \mathbf{Y} \quad (\textit{ s } \text{ is a t-conorm})$$

Intersection:  $R = P \cap Q$

$$R(x,y) = P(x,y) \textit{ t } Q(x,y) \quad \forall (x,y) \in \mathbf{X} \times \mathbf{Y} \quad (\textit{ t } \text{ is a t-norm})$$

Fuzzy relation  $R : \mathbf{X} \times \mathbf{Y} \rightarrow [0,1]$

Standard complement:  $\bar{R}$

$$\bar{R}(x,y) = 1 - R(x,y) \quad \forall (x,y) \in \mathbf{X} \times \mathbf{Y}$$

Transpose:  $R^T$

$$R^T(y,x) = R(x,y) \quad \forall (x,y) \in \mathbf{X} \times \mathbf{Y}$$



## **6.5 Cartesian product, projections, and cylindrical extension of fuzzy sets**

## Cartesian product

$A_1, A_2, \dots, A_n$  fuzzy sets on  $\mathbf{X}_1, \mathbf{X}_2, \dots, \mathbf{X}_n$

$$R = A_1 \times A_2 \times \dots \times A_n$$

$$R(x_1, x_2, \dots, x_n) = \min \{A_1(x_1), A_2(x_2), \dots, A_n(x_n)\} \quad \forall (x_i, y_i) \in \mathbf{X}_i \times \mathbf{Y}_i$$

## Generalization

$$R(x_1, x_2, \dots, x_n) = A_1(x_1) \, t \, A_2(x_2) \, t \, \dots \, t \, A_n(x_n) \quad \forall (x_i, y_i) \in \mathbf{X}_i \times \mathbf{Y}_i$$


$t = t$ -norm

# Examples


$$A(x) = \exp[-2(x - 5)^2]$$

$$B(y) = \exp[-2(y - 5)^2]$$

$$R = A \times B$$



$$R(x,y) = \min \{A(x), B(y)\}$$



$$R(x,y) = A(x)B(y)$$

# Projections of fuzzy relations

$$R: \mathbf{X}_1 \times \mathbf{X}_2 \times \dots \times \mathbf{X}_n \rightarrow [0, 1]$$

$$\mathbf{X} = \mathbf{X}_i \times \mathbf{X}_j \times \dots \times \mathbf{X}_k$$


$$R_{\mathbf{X}}(x_i, x_j, \dots, x_k) = Proj_{\mathbf{X}} R(x_1, x_2, \dots, x_n) = \sup_{x_t, x_u, \dots, x_v} R(x_1, x_2, \dots, x_n)$$

$$I = \{i, j, \dots, k\}, \quad J = \{t, u, \dots, v\}, \quad I \cup J = N, \quad I \cap J = \emptyset$$

$$N = \{1, 2, \dots, n\}$$

# Example

$$R(x, y) = \exp\{-\alpha[(x - 4)^2 + (y - 5)^2]\}, \quad \alpha = 1$$




$$R_X(x) = \text{Proj}_X R(x, y) = \sup_y R(x, y)$$

$$R_Y(y) = \text{Proj}_Y R(x, y) = \sup_x R(x, y)$$

# Example

$$R: \mathbf{X} \times \mathbf{Y} \rightarrow [0, 1], \quad \mathbf{X} = \{1, 2, 3\}, \quad \mathbf{Y} = \{1, 2, 3, 4, 5\}$$

$$R(x, y) = \begin{bmatrix} 1.0 & 0.6 & 0.8 & 0.5 & 0.2 \\ 0.6 & 0.8 & 1.0 & 0.2 & 0.9 \\ 0.8 & 0.6 & 0.8 & 0.3 & 0.9 \end{bmatrix}$$





$$R_X = [1.0, 1.0, 0.9]$$


$$R_Y = [1.0, 0.8, 1.0, 0.5, 0.9]$$

# Cylindrical extension


$$\text{cyl}A(x,y) = A(x), \quad \forall x \in X$$




$cylA$




$R$



$cylA \cup R$



$cylA \cap R$













## 6.6 Reconstruction of fuzzy relations

# Reconstruction using Cartesian product

$$\text{Proj}_X R \times \text{Proj}_Y R \supseteq R$$



$R$   
noninteractive



$R$   
interactive

## 6.7 Binary fuzzy relations

Binary fuzzy relation  $R : \mathbf{X} \times \mathbf{X} \rightarrow [0,1]$

## Features

### (a) Reflexivity


$$R(x,x) = 1$$

$$R(x,x) \supseteq I$$

$I = \text{Identity}$

$$R(x,x) \geq \varepsilon \quad \varepsilon\text{-reflexive}$$


$$\max \{R(x,y), R(y,x)\} \leq R(x,x) \quad \text{locally reflexive}$$



## (b) Symmetry


$$R(x,y) = R(y,x) \quad \forall x, y$$

$$R^T = R$$



## (c) Transitivity

$$\sup_{z \in \mathbf{X}} \{R(x, z) \wedge R(z, y)\} \leq R(x, y) \quad \forall x, y, z \in \mathbf{X}$$



# Transitive closure

$$\text{trans}(R) = \overleftrightarrow{R} = R \cup R^2 \cup \dots \cup R^n$$

$$R^2 = R \circ R \quad \dots \quad R^p = R \circ R^{p-1}$$

$$R \circ R(x, y) = \max_z \{ R(x, z) \wedge R(z, y) \}$$

If  $R$  is reflexive, then  $I \subseteq R \subseteq R^2 \subseteq \dots \subseteq R^{n-1} = R^n$

$I$  = identity

# Floyd-Warshall procedure to find $\text{trans}(R)$

**procedure** TRANSITIVE-CLOSUR-W ( $R$ ) **returns** transitive fuzzy relation

**static:** fuzzy relation  $R = [r_{ij}]$

**for**  $i = 1:n$  **do**

**for**  $j = 1:n$  **do**

**for**  $k = 1:n$  **do**

$\overleftrightarrow{r}_{jk} \leftarrow \max(r_{jk}, r_{ji} \text{ t } r_{ik})$

**return**  $R$



# Equivalence relations

$$R : \mathbf{X} \times \mathbf{X} \rightarrow \{0,1\}$$

$R$  is an equivalence relation if it is

- reflexive
- symmetric
- transitive

equivalence relations  
generalize the idea of  
equality

Equivalence class

$$A_x = \{y \in \mathbf{X} \mid R(x,y) = 1\}$$

$\mathbf{X}/R$  = family of all equivalence classes of  $R$  (partition of  $\mathbf{X}$ )

# Similarity relations

$$R : \mathbf{X} \times \mathbf{X} \rightarrow [0,1]$$

$R$  is a similarity relation if it is

- reflexive
- symmetric
- transitive

Equivalence class

$$P(R) = \{\mathbf{X}/R_\alpha \mid \alpha \in [0, 1]\}$$


Nested partitions: if  $\alpha > \beta$  then  $\mathbf{X}/R_\alpha$  finer than  $\mathbf{X}/R_\beta$

# Example

$$R = \begin{bmatrix} 1.0 & 0.8 & 0 & 0 & 0 \\ 0.8 & 1.0 & 0 & 0 & 0 \\ 0 & 0 & 1.0 & 0.9 & 0.5 \\ 0 & 0 & 0.9 & 1.0 & 0.5 \\ 0 & 0 & 0.5 & 0.5 & 1.0 \end{bmatrix}$$

$$R_{0.5} = \begin{bmatrix} 1 & 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 & 1 \end{bmatrix}, \quad R_{0.8} = \begin{bmatrix} 1 & 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}, \quad R_{0.9} = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1.0 & 1 & 0 \\ 0 & 0 & 1 & 1.0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

# Partition tree induced by similarity relation $R$



$$R_{0.5} = \begin{bmatrix} 1 & 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 & 1 \end{bmatrix}, \quad R_{0.8} = \begin{bmatrix} 1 & 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}, \quad R_{0.9} = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1.0 & 1 & 0 \\ 0 & 0 & 1 & 1.0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

# Compatibility relations

$$R : \mathbf{X} \times \mathbf{X} \rightarrow \{0,1\}$$

$R$  is a compatibility relation if it is

- reflexive
- symmetric

$\alpha$  -Compatibility class:  $A \subset \mathbf{X}$  such that

$$R(x,y) = 1 \quad \forall x,y \in A$$

Do not necessarily induce partitions

# Proximity relations

$$R : \mathbf{X} \times \mathbf{X} \rightarrow [0,1]$$

$R$  is a proximity relation if it is

- reflexive
- symmetric

Compatibility class:  $A \subset \mathbf{X}$  such that

$$R(x,y) = 1 \quad \forall x,y \in A$$

Do not necessarily induce partitions

$$R = \begin{bmatrix} 1.0 & 0.7 & 0 & 0 & .6 \\ 0.7 & 1.0 & 0.6 & 0 & 0 \\ 0 & 0.6 & 1.0 & 0.7 & 0.4 \\ 0 & 0 & 0.7 & 1.0 & 0.5 \\ 0.6 & 0 & 0.4 & 0.5 & 1.0 \end{bmatrix}$$